

A Solution to the μ Problem in Gauge-mediated Supersymmetry-breaking Models

T. Yanagida

Department of Physics, University of Tokyo, Tokyo 113, Japan

Abstract

We point out that a sector required to set the cosmological constant to zero in gauge-mediated supersymmetry-breaking models naturally produces a supersymmetry-invariant mass (μ term) for Higgs doublets of the order of the electroweak scale. Since this new sector preserves the supersymmetry, it does not generate supersymmetry-breaking masses for the Higgs doublets and thus the μ problem is solved.

Low-energy supersymmetry (SUSY) breaking with gauge mediation [1] has, recently, attracted much attention in particle physics. This is not only because it has a number of phenomenological virtues [2] but also because various mechanisms for dynamical SUSY breaking have been discovered [3][4]. This approach, however, has several drawbacks. The most serious one is the so-called μ problem. That is, if one wants to generate dynamically a SUSY-invariant mass (μ term) for the Higgs doublets by introducing their coupling to the SUSY-breaking sector, one necessarily gets too large SUSY-breaking mass (called B term) for the Higgs doublets [5]. Various solutions to this problem have been proposed [5] [1], but no compelling one has been found so far.

In this short paper we point out that a sector required to set the cosmological constant to zero in gauge-mediated SUSY-breaking (GMSB) models provides a natural solution to the μ problem.

Let us first discuss the cosmological constant problem in the GMSB models. In these models the SUSY is supposed to break down dynamically through nonperturbative effects of underlying gauge interactions. Vacuum-expectation values of the SUSY-breaking F term and the superpotential W are given by the dynamical scale Λ_{SUSY} of the underlying gauge interactions as

$$\langle F \rangle \sim \Lambda_{SUSY}^2 \quad (1)$$

and

$$\langle W \rangle \sim \Lambda_{SUSY}^3. \quad (2)$$

In supergravity the vacuum-energy density is fixed roughly by the vacuum-expectation values $\langle F \rangle$ and $\langle W \rangle$ as

$$\langle V \rangle \simeq \langle F \rangle^2 - \frac{1}{3M^2} \langle W \rangle^2, \quad (3)$$

where M is the gravitational scale $M \cong 2.4 \times 10^{18}$ GeV. Since the dynamical scale Λ_{SUSY} is much smaller than the M , the two terms in Eq.(3) never cancel out. Thus, we always have a positive vacuum-energy density (cosmological constant).

One may solve this problem by adding a constant term to the superpotential W by hand. However, it breaks a R symmetry. Once one allows explicit breaking of the R symmetry, there is no reason why the constant term should be so small compared to the gravitational scale M^3 . Therefore, it is quite reasonable to consider that such a constant term in the superpotential has a dynamical origin. Namely, we consider that the R symmetry is broken by nonperturbative effects of new gauge interactions so that the induced constant term in W cancels out the vacuum-energy density given by the SUSY-breaking sector. This requires fine tuning, but it seems the least assumption in the GMSB models.

We now construct a new sector for the R-symmetry breaking. The simplest example is a SUSY $SU(2)$ gauge theory with four doublet chiral superfields Q_i ($i = 1, \dots, 4$). We introduce six singlet superfields $S^{ij} = -S^{ji}$, where $i, j = 1, \dots, 4$. To make our analysis simpler we impose a global $SP(2)$ symmetry under which the Q_i transform as **4** and the S^{ij} split up as S_5^{ij} and S_0 transforming as **5** and **1**, respectively. We consider a continuous R symmetry $U(1)_R$ and assume the R-charges for all matter superfields Q_i and S^{ij} are $2/3$. Then a tree-level superpotential is given by*

$$W_{tree} = \lambda_5 Q_i Q_j S_5^{ij} + \frac{\lambda_0}{2} (Q_1 Q_2 + Q_3 Q_4) S_0 - \frac{g_5}{2} (S_5^{ij})^2 S_0 - \frac{g_0}{6} S_0^3. \quad (4)$$

It is known that the full nonperturbative effects in this theory generate a low-energy superpotential[6]

$$W_{dyn} = X(Pf M_{ij} - \Lambda^4), \quad (5)$$

where M_{ij} are the gauge-invariant degrees of freedom

$$M_{ij} = -M_{ji} \sim Q_i Q_j \quad (6)$$

and X is a Lagrange multiplier field. Then, the full low-energy superpotential is written in terms of the gauge invariants M_{ij} and S^{ij} as

$$W = X(Pf M_{ij} - \Lambda^4) + \lambda_5 M_{ij}^5 S_5^{ij} + \frac{\lambda_0}{2} (M_{12} + M_{34}) S_0 - \frac{g_5}{2} (S_5^{ij})^2 S_0 - \frac{g_0}{6} S_0^3. \quad (7)$$

*The model with $g_5 = g_0 = 0$ has been proposed as an example for the dynamical SUSY breaking in Ref.[3].

Here, M_{ij}^5 denotes the **5** representation of the $SP(2)$ while $(M_{12} + M_{34})$ transforms as a singlet under the global $SP(2)$.

This model possesses various SUSY-invariant vacua[†], among which we choose the following $SP(2)$ -invariant one[‡]:

$$\langle M_{12} \rangle = \langle M_{34} \rangle = \Lambda^2, \quad (8)$$

$$\langle S_0 \rangle = \sqrt{\frac{2\lambda_0}{g_0}} \Lambda, \quad (9)$$

$$\langle M_{ij}^5 \rangle = \langle S_5^{ij} \rangle = 0. \quad (10)$$

In this vacuum the continuous R symmetry breaks down to a discrete Z_{2R} , which generates a constant term in the superpotential W as

$$\langle W \rangle = \sqrt{\frac{8\lambda_0^3}{9g_0}} \Lambda^3. \quad (11)$$

Notice that there appears no R-axion superfield. This is because the R symmetry defined at the classical level is broken explicitly by nonperturbative effects at the quantum level since the R-current has an $SU(2)$ gauge anomaly.

Since the SUSY is preserved in this sector, the constant superpotential Eq.(11) leads to a negative vacuum-energy density

$$\langle V' \rangle \simeq -\frac{1}{3M^2} \langle W \rangle^2, \quad (12)$$

which is supposed to cancel the positive vacuum-energy density Eq.(3) coming from the SUSY-breaking sector. This gives us a relation between the two dynamical scales $\sqrt{\langle F \rangle} \sim \Lambda_{SUSY}$ and Λ as

$$\langle F \rangle^2 \simeq \frac{1}{3M^2} \langle W \rangle^2 \quad (13)$$

$$= \frac{8\lambda_0^3}{27g_0 M^2} \Lambda^6. \quad (14)$$

[†]In the limit of both g_5 and g_0 vanishing all the SUSY-invariant vacua run away to infinity [3].

[‡]In the $SP(2)$ -violating vacua we have Nambu-Goldstone superfields. If one introduces explicit breaking of the global $SP(2)$ one can eliminate such unwanted massless fields. In this case one may choose the $SP(2)$ -violating vacua, which does not, however, change any essential point in this paper.

We are now at the point to discuss the μ term for a pair of Higgs doublets H and $\overline{H}$. The form of coupling of the Higgs doublets to this R-symmetry breaking sector depends on the R-charge of the product $H\overline{H}$. We assume the product $H\overline{H}$ has the R-charge = 2/3. Then, the Higgs doublets have the following nonrenormalizable couplings to Q_i and S^{ij} in the superpotential:

$$W_{H,\overline{H}} = \frac{h}{2M}(Q_1Q_2 + Q_3Q_4)H\overline{H} + \frac{k}{M}(S_5^{ij})^2H\overline{H} + \frac{k'}{M}S_0^2H\overline{H}. \quad (15)$$

In the vacuum Eqs.(8)-(10) we have a SUSY-invariant mass μ for the Higgs doublets as

$$\mu = \frac{h}{2M}(\langle M_{12} \rangle + \langle M_{34} \rangle) = \frac{h}{M}\Lambda^2. \quad (16)$$

Here, we have neglected the contribution from the third term in Eq.(15) for simplicity, since it does not change the essential point of our conclusion. From Eqs.(14) and (16) we get

$$\mu \simeq \frac{3hg_0^{1/3}}{2\lambda_0} \left(\frac{\sqrt{F}}{10^6 GeV} \right)^{4/3} \times 100 GeV. \quad (17)$$

We see that the mass obtained for the Higgs doublets lies in the electroweak mass region for a reasonable range of parameters ($hg_0^{1/3}/\lambda_0 \simeq 0.1 - 10$ and $\sqrt{F} \simeq 10^5 - 10^7$ GeV [1][7]. Since the R-symmetry breaking sector preserves the SUSY, it never generates SUSY-breaking masses for the Higgs doublets. Thus, we have no μ problem.

It is now clear that the gauge invariant Q_iQ_j which condenses in the vacuum play a central role in our solution. We may construct various, similar models for the R-symmetry breaking sector. For example, consider an $SU(N_C)$ gauge theory with a pair of matter superfields Q and $\overline{Q}$ transforming as $\mathbf{N}_C$ and $\mathbf{N}_C^*$ under the $SU(N_C)$, respectively. Introduction of a singlet S allows us to have a tree-level superpotential

$$W_{Tree} = \lambda Q\overline{Q}S - \frac{g}{3}S^3. \quad (18)$$

In this model ($N_C > N_F = 1$) the full nonperturbative effects yield the exact effective superpotential at low energies as [8]

$$W_{eff.} = b\Lambda^{\frac{3N_C-1}{N_C-1}}(Q\overline{Q})^{\frac{-1}{N_C-1}} + W_{Tree}. \quad (19)$$

We have a SUSY-invariant vacuum

$$\langle Q\overline{Q} \rangle = \left(\frac{g}{\lambda^3} \frac{b^2}{(N_C - 1)^2} \right)^{\frac{N_C - 1}{3N_C - 1}} \Lambda^2 \quad (20)$$

$$\langle S \rangle = \frac{1}{\lambda} \left(\frac{g}{\lambda^3} \right)^{\frac{-N_C}{3N_C - 1}} \left(\frac{b}{N_C - 1} \right)^{\frac{N_C - 1}{3N_C - 1}} \Lambda. \quad (21)$$

As in the previous model a nonrenormalizable interaction $\frac{h}{M} Q\overline{Q}H\overline{H}$ produces the μ term for the Higgs doublets of the order of the electroweak scale.

In conclusion, we should stress that the vanishing cosmological constant in the GMSB models suggests the existence of new dynamics at the scale $\Lambda \sim 10^{10}$ GeV. We have no explanation for why the contributions from the SUSY-breaking and the R-symmetry breaking sectors to the vacuum-energy density cancel out exactly, but it seems a necessary condition for any realistic GMSB model. On the other hand, it is very encouraging that the new sector required to set the cosmological constant to zero naturally produces a μ term for the Higgs doublets in the desired mass region, $\mu \sim O(100)$ GeV.

Acknowledgment

We thank E. Stewart for useful comments.

References

- [1] M. Dine and A.E. Nelson, *Phys. Rev.* D48 (1993) 1277;
M. Dine, A.E. Nelson and Y. Shirman, *Phys. Rev.* D51 (1995) 1362;
M. Dine, A.E. Nelson, Y. Nir and Y. Shirman, *Phys. Rev.* D53 (1996) 2658.
- [2] For a review, see M. Dine, hep-ph/9612389.
- [3] K.-I. Izawa and T. Yanagida, hep-th/9602180, *Prog. Theor. Phys.* 95 (1996) 829.
K. Intriligator and S. Thomas, hep-th/9603158, *Nucl. Phys.* B473 (1996) 121.
- [4] K. Intriligator, N. Seiberg and S.H. Shenker, *Phys. Lett.* B342 (1995) 152;
P. Pouliot and M.J. Strassler, *Phys. Lett.* B375 (1996) 175;
C. Csaki, L. Randall and W. Skiba, hep-th/9605108;
Y. Shirman, hep-th/9608147.
- [5] G. Dvali, G.F. Giudice and A. Pomaral, hep-ph/9603238.
- [6] N. Seiberg, *Phys. Rev.* D49 (1994) 6857.
- [7] M. Hotta, K.-I. Izawa and T. Yanagida, hep-ph/9608147, *Phys. Rev.* D55 (1997) 415.
- [8] I. Affleck, M. Dine and N. Seiberg, *Nucl. Phys.* B241 (1984) 493.